

TOXICITY:

Harmful - This product may be harmful if swallowed. May cause organ damage from repeated oral exposure at high doses.

ECOTOXICITY

Designed for biocidal action.

Toxic to rodents and other terrestrial vertebrates if swallowed.

RODENTHOR GEL Rodenticide must not be applied onto or into water.

RODENTHOR GEL Rodenticide must be applied as a contained, ground-based application only.

PRECAUTIONS

Keep out of reach of children. Do NOT touch bait with bare hands. Do NOT place bait in the open or places accessible to children, domestic pets, animals or birds. After use, wash hands and exposed skin thoroughly with soap and water. Avoid contact with food, food utensils or places where food is prepared or stored.

DIRECTIONS FOR USE**RESTRAINTS**

DO NOT apply bait to crops.
DO NOT place bait in the open.
DO NOT exceed 3 m between bait stations for mice or 9 m for rats in Clean-Out treatments.

Baiting Strategy

- Eliminate as far as practical, all alternative food sources.
- Bait infested area. Place baits only within tamper proof bait stations.
- Actively search for, remove and destroy dead and moribund rodents whilst bait is being used.
- Ensure baits are removed when rodent activity has ceased.

Clean-Out Treatments (Pulse Baiting)

- Clean-Out treatments are used to remove existing infestations of rats and mice.
- Always record the number and locations of bait stations for the purpose of bait replacement and later removal.
- For rats, apply 15 g – 30 g bait gel to the base of bait stations at intervals of 5 to 9 m in infested areas
- For mice, apply 3 to 4 g bait gel to the base of bait stations at 2 to 3 m intervals in infested areas.
- Place baits so as to prevent access to non-target animals. Where access to non-target animals cannot otherwise be prevented, secured tamper resistant bait stations must be used.
- Use the greater number of baits and shorter distance between bait placements when infestations are heavy.
- Conduct initial inspection of bait points after 3 or 4 days. Replace any bait that has been consumed. Repeat approximately 4 days later and then weekly. Replace contaminated or spoiled baits.
- Baiting for at least 2 weeks (3 to 4 pulses) will be necessary to reduce rat/mouse numbers to a low level. Although heavy infestations may require longer treatments to achieve complete eradication.
- Discontinue the treatment when effective control has been achieved and remove all bait and bait stations unless a maintenance treatment is to be undertaken.

Monitoring and/or Maintenance Treatments

- Monitoring and maintenance treatments are required at sensitive sites where there is no current rodent infestation but where the prevention of such infestation is essential to prevent significant risks to human health. Such treatment may be a condition for audit approval within food safety guidelines. However, all use of this product must comply with the conditions of this label.
- Always record the number and locations of bait stations for the purpose of bait replacement and later removal.
- For rats and mice apply up to 15 g bait gel to bait stations at intervals of between 15 and 30 m in areas prone to rodent infestation.
- Place baits so as to prevent access to non-target animals. Always use tamper-resistant bait stations for monitoring and maintenance treatments and fix the stations securely to the substrate.



Batch No.:

Date of Manufacture:

HARMFUL
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**ACTIVE INGREDIENT:**

**Contains 0.05 g/kg brodifacoum
in the form of a ready to use bait**

**For effective control of RATS and
MICE around industrial, commercial,
agricultural and residential buildings.
Controls rodents resistant to other
anticoagulants, including warfarin,
coumatetralyl and bromadiolone
ONE FEED KILLS RATS AND MICE***

NET CONTENTS:

300 g

Manufactured by:

Zapi SpA

Via Terza Strada 12, 35026 Conselve, Padua, Italy

NZ Agent:

Ensysstex New Zealand Ltd

17C Corinthian Drive, Albany 0752, Auckland

www.ensystex.co.nz

CUSTOMER SERVICE 0800 ENSYSTEX (0800 367 978)

Registered pursuant to the ACVM Act 1997, No V9666
See www.foodsafety.govt.nz for registration conditions

Approved pursuant to the HSNO Act 1996, Approval Code HSR101434
See www.epa.govt.nz for approval controls

*Registered trademarks of Ensysstex, Inc. used by licence to Ensysstex New Zealand Ltd.

- Bait stations should be checked at regular intervals according to an agreed and documented schedule.
- Use placebo or unpoisoned baits if the placement of the bait away from buildings presents a hazard to non-target animals.
- When a rodent infestation is detected the operator should revert to the directions for Clean-Out treatments.

Application

| SITUATION | PESTS | RATE | CRITICAL COM-MENTS |
|------------------------------------------------------------------------------|--------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------|
| Industrial, Commercial, Public Services, Agricultural and Domestic Buildings | Rats (<i>Rattus rattus</i> , <i>R. norvegicus</i>) Mice (<i>Mus musculus</i>) | 15 g to 30 g (15 to 30 mL) bait per bait point 3 to 4 g (3 to 4 mL) bait per bait point | Set out bait in places frequented by mice and rats in and around buildings or enclosed spaces, e.g. drains. |

GENERAL INSTRUCTIONS

This product may only be used according to this label. DO NOT place RODENTHOR LIPO-GEL out in the open. Ensure bait is secure and inaccessible to children, pets, domestic animals and wildlife. Place bait along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Record the number and location of bait placements for the purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait. Look for dead rats or mice, burn or bury them. DO NOT place in refuse bins or rubbish tips. Ensure baits are removed when rodent activity has ceased.

Note: One feed is sufficient to kill rats and mice. Although rats and mice will continue to eat the bait after a lethal dose has been eaten, the directions for use are intended to minimise this. Insert the tube into the caulking gun and cut a hole in the conical tip of the tube and screw on the nozzle. Cut the nozzle above the screw thread taking care not to damage the thread itself, then dispense the product. Once finished, replace the protective cap and release the pressure of the caulking gun to prevent loss. Keep the tube upright.

Depending on the amount eaten, this bait may kill or injure. The bait is hazardous to dogs, cats, pigs, poultry and other wildlife. DO NOT contaminate dams, waterways and drains with the bait or its used container.

STORAGE AND DISPOSAL:

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Do not burn empty containers or product. DO NOT use containers which have held bait for any other purpose. Dispose of carcasses safely by burning or burying in an approved landfill.

SAFETY DIRECTIONS

Poisonous if swallowed. DO NOT touch bait. If on skin, and after each baiting, wash thoroughly with soap and water. Wash hands after use.

FIRST AID

If swallowed, do not induce vomiting. For advice contact the National Poisons Centre. 0800 POISON (0800 764 766) or a doctor immediately. Vitamin K1 (Phytomenadione) is antidotal.

NOTE TO PHYSICIANS AND VETERINARIANS

Brodifacoum, the active constituent of RODENTHOR LIPO-GEL Rodenticide, is a long-lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet and is available from Ensysstex on request or visit our web site at www.ensystex.co.nz

NOTICE

Ensysstex New Zealand Ltd warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions for use under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of the product contrary to label instructions or under abnormal conditions.

L-NZ-RDTG1002 1.02 01.22

ENSYSTEX